
Heartworm/4Dx Testing



Fecal Exams/Deworming

Date	Pos	Neg	Notes

Medical/Surgical History

Date	Notes
	-

Rabies Tag Information

Date	Number



			Zip	
	Sex		State	

Telephone

Canine Health Record:

Date of Birth

Markings

Breed

Owner Address

Vaccination History

Date	Distemper	Adenovirus	Parainfluenza	Parvovirus	Lepto	Coronoa	Lyme	Bordetella	Rabies

Vaccination

Rabies

Of all animal diseases, rabies is the most feared. Most pets are exposed to rabies by bites from wild animals, particularly skunks, bats, raccoons, and foxes. The disease can be transmitted to humans through the bite or scratch of an infected animal, and is nearly always fatal. Vaccination of all dogs and cats is the most effective means of prevention.

Canine Distemper

This highly contagious viral disease is found wherever dogs are found. It affects the respiratory and nervous system and is often fatal. Primary vaccination should begin at 6-8 weeks as the disease can be contracted at a very young age.

Canine Parvovirus (CPV)

This common contagious viral disease causes severe diarrhea and vomiting in dogs of all ages, but is deadly in puppies.

Canine Coronavirus (CCV)

Coronoavirus is highly contagious and causes a milder form of gastrointestinal symptoms than parvovirus. We do not include it in our core vaccinations.

Canine Parainfluenza

This viral respiratory disease is often partly responsible for "kennel cough" in dogs. Infection can be severe in young puppies. Parainfluenza protection is often included in combination distemper-parvovirus vaccinations.

Canine Adenovirus Type 1 & 2

Canine Adenovirus Type 1 infection causes infectious hepatitis, which can lead to kidney damage. Type 2 can be a complicating factor in kennel cough.

Canine Leptospirosis

Leptospirosis is a bacterial disease carried by many wild animals. A dog can contract the disease from infected animals by drinking contaminated water. Yearly vaccination decreases the odds of your dog acquiring the disease.

Canine Bordetella

Frequently involved in kennel cough, this bacterial infection may occur simultaneously with distemper, adenovirus type 2, parainfluenza, and other respiratory infections.

Lyme Disease

Borreliosis, or Lyme Disease, is an infection caused by a bacteria which is spread through the painless bite of an infected tick. Symptoms include fever, lethargy, muscle stiffness, joint pain, and loss of appetite.

Grand Avenue Veterinary Clinic

5503 Grand Avenue Duluth, MN 55807 Phone: (218) 628-0301 Fax: (218) 628-1448 www.grandavevetclinic.com